

Land use patterns and influences of protected areas on mangroves of the eastern tropical Pacific

Juliana López-Angarita^{a,b,*}, Alexander Tilley^{b,c}, Julie P. Hawkins^a, Carlos Pedraza^d, Callum M. Roberts^a

^a Environment Department, University of York, York YO10 5DD, United Kingdom

^b Fundación Talking Oceans, Carrera 16 # 127 – 81, Bogotá 111831, Colombia

^c WorldFish, Ministerio de Agricultura e Pesca, Dili, Timor-Leste

^d Faculty of Natural Sciences and Mathematics, Universidad del Rosario, Carrera 26 No 63B-48, Bogotá, Colombia

ARTICLE INFO

Keywords:

Agriculture
Coastal development
Aquaculture
Land-cover change
Land-use change
Wetland

ABSTRACT

Mangroves are one of the most productive ecosystems in the world, sustaining millions of coastal livelihoods. However, their area of occurrence has been greatly reduced over the last century. In this study, we identify potential drivers of land use and land cover change adjacent to mangroves on the Pacific shorelines of Colombia, Panama and Costa Rica. We also evaluate the effectiveness of protected areas at halting mangrove deforestation between 2000 and 2012. Across all countries, agriculture was the most dominant land use type adjacent to mangroves, inside and outside protected areas. Results show that a combined total of 564 ha were lost, representing an average loss rate of only 0.02% per year. 75% of the total mangrove loss occurred in locations outside protected areas, with only 138 ha cleared from inside protected areas. Results suggest current conservation policies for mangrove protection in the study countries are effective at reducing deforestation and set a positive example for regions where mangroves are in decline.

1. Introduction

It is estimated that by 2050, global crop production must double to meet the demands of a rising global population (Tilman et al., 2011). Despite suggestions to prevent the increase of cultivated area, the global pattern of increasing agricultural field sizes is often driven by government incentives, demand for biofuels, and technology (White and Roy, 2015). Worldwide rates of urban land expansion are higher than, or equal to, urban population growth rates (Seto et al., 2011). It is therefore expected that Land Use and Land Cover Change (LULCC) will increase as global population grows and developing countries become more affluent.

As LULCC intensifies, the effects of arable and urban land expansion may have significant and potentially irreversible consequences on ecosystem function and integrity (Foley et al., 2005). For instance, land conversion that removes primary forest has been shown to greatly reduce species diversity (Gibson et al., 2011). In the tropics, LULCC is associated with agricultural products for food, feed, and fuel (Gibbs et al., 2010; Blanco et al., 2012). Human reliance on natural environments is high in these regions and more than half of the new agricultural land created between 1980 and 2000 was via deforestation

(Gibbs et al., 2010).

Mangrove forests are restricted to the interface between land and sea in tropical and subtropical latitudes. They are highly productive, provide a vast array of ecosystem services (Hogarth, 2007), and diversify and sustain livelihoods for millions of people (UNEP, 2014). Despite these widely appreciated values, mangrove cover is rapidly declining in different regions (Valiela et al., 2001; Alongi, 2008; Richards and Friess, 2015).

Estimates of global mangrove loss vary across regions and between methods used (Alongi, 2002; Giri et al., 2011; López Angarita et al., 2016). The development of optical remote sensing technology has allowed for a better estimation of mangrove coverage, and for the exploration of LULCC dynamics (Manson et al., 2001; Dahdouh-Guebas et al., 2004). Recently, development of new radar technology sensitive to forest spatial structure has allowed for accurate estimates of mangrove deforestation rates (Lucas et al., 2007; Simard et al., 2008; White and Roy, 2015; Hamilton, 2013; Thomas et al., 2017). However to date there is little information on the proximate drivers of LULCC in mangrove forests or replacement land uses (Tilman et al., 2011; Richards and Friess, 2015).

The Eastern Tropical Pacific (ETP) biogeographical region spans the

* Corresponding author at: Fundación Talking Oceans, Carrera 16 # 127 – 81, Bogotá 111831, Colombia.

E-mail addresses: julianalop14@gmail.com (J. López-Angarita), carlosa.pedraza@urosario.edu.co (C. Pedraza).



Fig. 1. Geographical extent of the study (red line), on the Pacific coasts of Costa Rica, Panama, and Colombia (shaded green). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

continental shelf and oceanic islands of Southern Baja California to northern Perú (Briggs, 1974), and supports a range of rich fisheries and exhibits many endemic species (Zapata and Robertson, 2006; Fiedler and Talley, 2006; Hogarth, 2007). In terms of mangrove protection in the region, 58% of mangroves that occur on the Pacific coast of Costa Rica are inside protected areas, compared to 51% in Panama and 28% in Colombia (López Angarita et al., 2016).

Mangrove cover in the ETP has followed global trends of decline, with its greatest loss occurring between the 1960s and 1990s (Valiela et al., 2001; López Angarita et al., 2016). Since then, countries in the ETP have strengthened their conservation policies for mangroves, via creation of protected areas and laws regulating mangrove use (Lacerda et al., 1993; ANAM-ARAP, 2013; López Angarita et al., 2016). To date there has been little or no assessment of the effectiveness of this protection.

In this study, we aim to identify the potential drivers of mangrove decline on the Pacific coasts of Costa Rica, Panama, and Colombia (Fig. 1) by mapping anthropogenic activities of LULCC in mangroves and performing analyses by country to compare trends within the region. Additionally, we determine the effectiveness of mangrove conservation policies by calculating rates of mangrove deforestation inside and outside protected areas, between 2000 and 2012.

2. Methods

2.1. Mangrove forest loss

To calculate the rate of mangrove deforestation we used the Global Forest Change dataset created by Hansen et al. (2013), which provides an index of annual deforestation between 2000 and 2015 per pixel (pixel size of 0.09 ha). These data are available up until 2015, but our study used the data between 2000 and 2012 only, to align with available land use data. We projected the Global Forest Change dataset for

each country using UTM 18 N/17 N transformed from WGS84. Country-level mangrove areas were identified by overlaying the political limits of the studied countries with the global distribution of mangroves (Mangrove Forests of the World) in 2000 provided by Giri et al. (2011). Offshore islands were not included in our study. In a small section of the Colombian Pacific coast, we found a projection error causing misalignment of mangroves with the coastline in the Giri et al. (2011) global dataset, so we used Google satellite imagery and the mangrove distribution dataset for Colombia (IDEAM et al., 2007) to correct the error by manually fitting mangrove area polygons to the coastline. These steps resulted in a data layer of mangrove deforestation by country and year for the region of interest. This layer was used to calculate the percentage of mangroves deforested in the region for each country using number of pixels to estimate area. We obtained the rate of deforestation per year by dividing the percentage lost by the 12-years sampled (2000 not included and 2012 included). We used the same input layers (Global Forest Change and Mangrove Forests of the World) that Hamilton and Casey (2016) used, with a different methodological approximation, in their Global Database of Continuous Mangrove Forest Cover for the 21st Century, so we could compare our results with their deforestation rates.

2.2. Potential drivers of LULCC in mangrove areas

Ten different datasets of land cover with a resolution $\leq 30 \text{ m}^2$ were used to map the distribution of potential drivers of LULCC across the three countries (Table A1). We grouped potential drivers into three major classes: aquaculture, agriculture (includes cattle farms, oil palm plantations, and crops such as rice and fruits), and coastal development. Coastal development included towns and infrastructure such as ports and agricultural processing plants. Infrastructure was not analyzed as a separate class due to the few records associated with it. In this manuscript we used the term “potential drivers” to define land use types with

the potential to negatively impact mangroves as shown elsewhere in the literature (e.g. Hamilton, 2013; Richards and Friess, 2015; Thomas et al., 2017). However, it is important to clarify that in the studied period, these current land uses might not have been directly responsible for the loss of mangroves in our study area.

To quantify the spatial distribution of potential drivers of LULCC adjacent to mangroves, we used an overlaid 1 km² grid to divide the study area into sample units. A 1 km² grid was selected to simplify interpretation of results at scale, as our model was designed to be a tool for managers. Grid squares were ground-truthed in all countries between 2013 and 2015 to calibrate the interpretation of land use in the datasets. We chose areas to ground-truth based on the significant presence of anthropogenic activities close to mangroves. Of 9812 1 km² grid cells placed over mangroves of the studied region, 401 were ground-truthed (see ground-truthing section in Appendices and Table A2). We aimed to ground-truth areas that were highly representative of the mangrove-land use landscape in each country, but for logistical reasons, it was not possible to visit the south of Colombia. Therefore, our interpretations might be biased for this area. When errors were present in the land use classification of the datasets, we used Google Earth images calibrated with ground-truthing to re-classify the polygons.

To display the spatial patterns of potential drivers of LULCC adjacent to mangroves, we developed a cumulative model, where grid cells were scored based on the presence (1) or absence (0) of aquaculture, agriculture, and coastal development throughout the region. Scores were summed per cell to return a possible value between 0 and 3. The proportion of cells belonging to the different values of cumulative scores was calculated. Due to the complexity of measuring the cascading effects that potential drivers of LULCC have in mangroves on a regional scale, the same weighting was applied to all potential drivers. Cells were given a color scale according to the total score. The extent of different land use types and the impact score results were compared between countries using Chi-square tests.

2.3. Protected areas

We compared the extent of mangrove deforestation inside and outside protected areas between 2000 and 2012 by mapping the boundaries of protected areas present on the Pacific coast of the countries studied according to government datasets (Tables A1, A3). We used the global forest change dataset (Hansen et al., 2013) to estimate figures of mangrove deforestation within protected areas. Protected areas established after the year 2000 were analyzed separately to accurately assess how deforestation had occurred inside and outside during the study period. Proportion of mangrove loss inside and outside protected areas was compared between countries using a Chi-square test. Finally, we compared the distribution of potential drivers of LULCC inside and outside protected areas by estimating the proportion of cells for each land use type. Cumulative score of mangroves was also compared inside and outside protected areas. Cells divided by protected area boundaries were classified as inside protected areas. Given the diverse types and protection levels of protected areas in the region, a comparative inside/outside approach was used. Geospatial analyses and calculations were performed in ArcGIS 10.3.1 and statistical tests in JMP version 13.

3. Results

3.1. Mangrove forest loss

The total area of mangroves on the Pacific coast of each country showed that Colombia has the largest area, followed by Panama then Costa Rica (Table 1). Over the reporting period, 564 ha or 0.18% of the total mangrove area were lost in all countries combined (Table 1). In Costa Rica by 2012, 0.32% of mangroves present in 2000 had been

deforested, with figures of 0.21% for Panama and 0.11% for Colombia.

Across all countries studied, the average annual deforestation rate was 0.02%. Temporal trends of mangrove deforestation showed that deforestation peaked in Panama and Costa Rica in 2008. An increasing trend of forest loss was observed in Colombia, whereas in Costa Rica deforestation has decreased with time (Fig. 2).

3.2. Potential drivers of LULCC in mangrove areas

According to the model, around 60% of cells across the three countries combined had no adjacent potential drivers of LULCC, whereas in 40% of cells, one or more land use types were present (Fig. 3). The proportion of cells with potential drivers was significantly different for three countries ($\chi^2(6, N = 9812) = 1132.36, p < 0.0001$), as well as the extent of different land use types, including mangroves ($\chi^2(4, N = 4058) = 712.45, p < 0.0001$). In Colombia, 73% of cells with mangroves had no potential drivers, whereas in ~26% of cells, one or more land use types were present (Table A4, Fig. A1). In this country, the most common land use was agriculture, present in 26% of cells within the grid, followed by coastal development (1.6%), while aquaculture did not occur (Fig. 4). In Panama, potential drivers were present in 53% of the cells with mangroves (Table A4). Agriculture was the most dominant land use, present in 30% of the cells, followed by coastal development (19%). 6% of cells were adjacent to aquaculture ponds (Fig. 4). In Costa Rica, 60% of mangrove cells had no proximate land use. Agriculture was the most common land use (28%), while aquaculture and coastal development had an equal representation of 9%.

3.3. Protected areas

Of the 31 protected areas mapped on the Pacific coast of Panama, 17 contained mangroves; in Colombia 6 of 9 contained mangroves; and for Costa Rica, 23 of 53 (Table A3). While figures for deforestation inside and outside protected areas varied between the three countries, significant differences between mangrove loss inside and outside were supported statistically for all countries ($\chi^2(2, N = 564) = 53.06, p < 0.001$), with loss inside protected areas lower than outside in all cases (Fig. 5). Across all three countries, 75% of deforestation occurred outside protected areas.

Ninety two percent of cells with mangroves inside protected areas in Colombia had no adjacent land use (Table A5). In Panama, agriculture was present in 50% of mangrove cells inside protected areas. In Costa Rica, 68% of mangrove cells inside protected areas had no proximate land use, 23% were surrounded by agriculture, and 8% by coastal development (Table A5).

4. Discussion

Knowledge of recent trends of mangrove deforestation is important in evaluating the effectiveness of current conservation policies. In a widely cited paper, Duke et al. (2007) raised concerns about the high rate of mangrove loss and estimated that the world could be without functional mangroves within 100 years. Fortunately, this scenario was based on extrapolated rates of mangrove deforestation from 1980s and 1990s and does not seem feasible now, as current research has shown that these trends appear not to have continued into the 21st century. This is certainly true in the ETP, where our calculated average annual loss rate of 0.02% between 2000 and 2012, confirm findings by Hamilton and Casey (2016), who report similar deforestation rates for the study countries (including the Caribbean coasts) (Table 1). Moreover, it is likely that our estimates are similar to country level estimates because the Pacific coast represents the majority of mangrove forest coverage for all study countries. Hamilton and Casey (2016) found a global rate of mangrove deforestation of 0.16%, with the highest levels of deforestation in Southeast Asia, particularly in Indonesia exhibiting an annual rate of 0.26%–0.66% per year. Our results support the low

Table 1

Figures for mangrove deforestation between 2000 and 2012 on the Pacific coasts of Costa Rica, Panama and Colombia. For comparison purposes, mangrove deforestation rates found by Hamilton and Casey (2016) using the Global Database of Continuous Mangrove Forest Cover for the 21st Century (CMGFC-21), are also shown.

Country	Mangrove area in the Pacific coast (ha)	Mangrove area lost since 2000 (ha)	% of total area lost 2000–2012	Annual loss rate (% of total area per year)	National annual loss rate (CMGFC-21)	% of total area lost 2000–2012 (CMGFC-21)
Costa Rica	37,266.5	120.4	0.32	0.03	0.029	0.35
Panama	135,955.8	287.7	0.21	0.02	0.025	0.29
Colombia	141,271.6	156.0	0.11	0.01	0.011	0.137
Total	314,493.8	564.1	0.18	0.02	NA	NA




Fig. 2. Temporal trends in the deforestation of mangroves between 2000 and 2012 on the Pacific coasts of Panama, Colombia, and Costa Rica. The lower right panel shows cumulative forest loss for all countries.

estimated deforestation rates in ETP countries, suggesting these low rates might be compensating for higher losses in other regions and thus, counterbalancing global mangrove loss rates.

In the ETP, annual loss rates calculated prior to 2000 were higher than 1% for the three countries included in this study, due to inclusion of figures of historic deforestation, when most mangroves were lost (Valiela et al., 2001; López Angarita et al., 2016). Recent trends of mangrove deforestation are consistent with historical trends, with Panama displaying the largest losses and Costa Rica the lowest (López Angarita et al., 2016). Despite Costa Rica having declared all mangroves no-take areas in 1998 (Valiela et al., 2001; RAMSAR, 2015), it showed the highest annual rate of loss in this study. Temporal patterns of deforestation illustrate that Costa Rica is the only country showing a declining trend in recent years, while Panama and Colombia exhibit gradual increases in deforestation rates.

The rates of deforestation in the study countries are low compared to post-2000 deforestation in other regions and forest types (Valiela et al., 2001; Potapov et al., 2012; Nepstad et al., 2014; Richards and Friess, 2015), which suggests that mangrove protection is relatively effective. In contrast, Richards and Friess (2015) estimated that between 2000 and 2010, mangroves in South East Asia were being lost at an average rate of 0.18% per year. For the same time period, in the

Democratic Republic of the Congo, average annual gross forest loss was 0.23% of forest area, across all forest types (Potapov et al., 2012). However, other studies have also reported significant declines in deforestation rates for other forests: In the Brazilian Amazon, forest loss declined by 70% between 2005 and 2013, passing from a ten year average of 19,500 km² per year, to 5843 km² (Nepstad et al., 2014). National deforestation rates, across all forest types, decreased after the year 2000 in Costa Rica, and Colombia (Cabrera et al., 2011). Perhaps due to its isolation, the Pacific coast region of Colombia has had the least amount of forest loss nationwide (Cabrera et al., 2011). Despite overall deforestation in Panama having decreased compared to the 1990s, figures remain quite high for the region, with an annual rate of 0.41% between 2000 and 2008 (Mariscal, 2012).

The cumulative model presented here offers perspective on the threats affecting mangroves at national scales, determined by proximity of potentially damaging activities adjacent to mangroves. Our analysis found that agriculture is now consistently the most dominant potential driver of LULCC adjacent to mangroves outside and inside protected areas. Ground-truthing showed that rice, watermelon, melon, sugar cane, and oil palm are the main crops grown, and that cattle farming also occurs. The intensity and extent of agriculture adjacent to mangroves varied among the countries examined, with small-scale




Fig. 3. Map of a cumulative model of potential drivers of land use and land cover change in mangroves on the Pacific coasts of Costa Rica, Panama and Colombia. Color grid represents 1 km² cells where the analysis was performed. Green cells represent mangroves without adjacent potential drivers of land use and land cover change, blue cells represent mangroves adjacent to one potential driver, yellow cells are mangroves adjacent to two land use types, and red cells are mangroves adjacent to three land use types. For visualization purposes only, red insets provide a magnified view of the selected area. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 4. Distribution of agriculture, aquaculture, and coastal development next to mangroves on the Pacific coasts of Colombia, Panama, and Costa Rica. Percentages are calculated from a 1 km² grid placed over the mangroves of the Pacific coast of each country.

agriculture prevalent on the Pacific coast of Colombia, in contrast to more productive agro-economic regions in Panama and Costa Rica (Pinto and Yee, 2011). In Panama, rice and beef are the most commercially important agricultural commodities, and they are produced in rotation on the same land (Trejos et al., 2008). In Costa Rica, melon and oil palm have the highest yield per hectare, are planted in high-density monocultures, and receive large inputs of chemical pesticides and fertilizers (Bach, 2007). Agriculture, particularly rice, has been shown to be an increasingly important driver of mangrove loss in other regions

also (Richards and Friess, 2015; Thomas et al., 2017).

Coastal development was a frequent potential driver of LULCC in Panama. In this country, mangroves adjacent to urban zones are commonly converted to areas of development for tourism and urban expansion (Benfield et al., 2005). On the contrary in Colombia, most of the Pacific coast population is scattered in small villages only accessible by boat (García, 2010). This isolation from the rest of the country translates into good coverage of natural rainforest and mangroves (Sánchez-Paez et al., 1997). However, this lack of infrastructure and




Fig. 5. Total mangrove deforestation in hectares between 2000 and 2012 (black bars), highlighting deforestation inside (grey bars) and outside (white bars) protected areas on the Pacific coasts of Colombia, Costa Rica, and Panama.

accessibility fosters poverty and drives a high dependence on natural resources (Leal, 2000; Blanco et al., 2011).

Aquaculture, in particular extensive shrimp aquaculture, is widely claimed to be the most important driver of mangrove loss worldwide (Páez-Osuna, 2001; Alongi, 2002; Giri et al., 2008, 2011). Yet, in the three countries studied, it was far less important than agriculture, and not even observed as a land use in the Colombian Pacific. The low importance seen in this region may be explained by large scale abandonment of ponds (Bolanos, 2012) following the outbreak of “white spot virus” in Central America, that affected both wild and cultured shrimp (Nunan et al., 2001).

Data used in this study to quantify forest loss were derived from Landsat images with a resolution of 30x30m. It is possible that the spatial resolution used underestimates mangrove deforestation by not detecting losses at smaller scales. Deforestation is likely to happen at the interface between forests and other land use types (Etter et al., 2006), which makes it hard to detect in satellite images (Heumann, 2011; Thompson et al., 2013). For example, in the Gulf of Montijo, Panama, it was reported that the area of rice crops adjacent to mangroves has increased gradually (ANAM, 2004) but the figures have not been quantified. Additionally, humid tropical regions such as the ETP are particularly challenging to map given the consistent cloud cover that affects the clarity of satellite images (Gibbs et al., 2010; Heumann, 2011). The ETP also shows large transitions between mangroves and other forests types, in some instances related to river discharge diversions (Restrepo and Cantera, 2013; Parra and Angel, 2014) where deforestation is likely to become cryptic. Therefore, there is underlying potential for underestimation of the integrity of mangroves, as satellite images at the scale used in this study do not allow for the identification of small-scale forest clearing and the slow rate of habitat degradation at forest fringes and mangrove transitions. This bias may be even more significant for Colombia, where ground-truthing was limited, as cryptic

Appendix A

Name and source of layers used for spatial analysis (Table A1); detail of ground-truthed areas (Table A2); list of protected areas included in this study (Table A3); cumulative model results (Table A4); cumulative model results inside and outside protected areas (Table A5); frequency distributions of cumulative model results (Fig. A1).

ecological degradation due to selective logging, and clear cutting of basin mangroves is known to occur (Blanco et al., 2012). Nevertheless, our approach uses robust data of forest loss (Hansen et al., 2013) that has been applied to quantify forest cover change in many other regions worldwide (Potapov et al., 2012; Hansen et al., 2013; Richards and Friess, 2015; Hamilton and Casey, 2016). It is important to highlight however, that primary data sources, radar, high-resolution aerial images, and field ecological assessments are highly needed to improve the estimation of deforestation rates in the challenging environment of the ETP (Blanco-Libreros and Estrada-Urrea, 2015). As such, rates of loss are likely to be higher than the estimates presented here for the above reasons.

Our analysis showed that most mangrove deforestation (75%) occurred outside protected areas. In Colombia, this equated to ~8 ha, whereas the figures for Costa Rica and Panama were 49 ha and 81 ha respectively. Other studies of tropical forests have also shown that the presence of protected areas significantly reduces deforestation inside them (Bruner, 2001; Naughton-Treves et al., 2005; Andam et al., 2008; Gaveau et al., 2009; Miteva et al., 2015; Spracklen et al., 2015), and this has been shown specifically for mangrove forests (Miteva et al., 2015). Despite protected areas in this region being often undermanaged (López Angarita et al., 2014) and under increasing stress from human activities (Chape et al., 2005), our findings provide reassuring evidence that despite the diverse management approaches represented in this analysis (Table A3), protection has had an overall positive effect in reducing mangrove deforestation. All study countries recognize in law that any activity intended to exploit or modify mangroves requires prior government evaluation and permission (García, 2010; Salas et al., 2012; ANAM-ARAP, 2013; López Angarita et al., 2016).

In the ETP, communities have been reported to have increasing participation in the management of mangroves (Kaufmann, 2012; Kothari et al., 2015; Vieira et al., 2016). Our results provide evidence that these initiatives combined with government input, are effective at reducing mangrove loss, and set a positive example for other regions where this ecosystem is being degraded. Our results are relevant to conservation and policy making in the region as they highlight the relative successes of formal protection of mangroves in the ETP. We recommend participatory land use planning at the community level to empower local stakeholders in mangrove protection.

Acknowledgements

We gratefully acknowledge the support (via fellowships awarded to J. López-Angarita) of the Colombian Department of Science and Technology COLCIENCIAS, the Schlumberger Foundation Faculty for the Future Programme, the World Wildlife Fund's Russell E. Train Education for Nature program (Agreement No. RU63), the Smithsonian Tropical Research Institute, and the Save our Seas Foundation (SOSF2013APP2N). We thank R. Venegas-Li, L. LeVay and R. Marchant for their comments on the manuscript. Special thanks to the governments of Colombia, Costa Rica, and Panama for sharing spatial data. Thanks to J. Cubillos and M. Villate for their help in the field.

Table A1

Details of the layers used to analyse proximate drivers of land use and land cover change adjacent to mangrove forest on the Pacific coastlines of Colombia, Costa Rica and Panamá.

Country	Variable	Name of layer and source
Colombia	Land use and land cover	National Cartographic database 2000–2009. National layer of land cover (CORINE land Cover) 2005–2009. Instituto Geográfico Agustín Codazzi.
	Mangroves	Continental, coastal and marine ecosystems of Colombia (IDEAM et al., 2007).
Costa Rica	Protected areas	Limites de áreas protegidas. Parques Nacionales Naturales de Colombia. 2014–2015.
	Land use and land cover	Atlas Nacional de Costa Rica, 2008. Global land cover - GlobeLand30 (Chen et al., 2015). www.globallandcover.com .
Panamá	Mangroves	Inventario Forestal de Costa Rica 2005.
	Protected areas	Sistema Nacional de Áreas de Conservación (SINAC) 2013.
	Land use and land cover	Land use Panamá 2008. Forest cover inventory 2000. Sistema Nacional de Áreas Protegidas (SINAP).

A.1. Ground truthing

Ground-truthing trips were made in 2013–2015 to all countries studied. Areas to ground-truth were chosen according to the significant presence of anthropogenic activities close to mangroves. Places visited in Costa Rica were: the Nicoya peninsula, Golfo Dulce and the Terraba Sierpe wetland; in Panamá: Aguadulce district, Gulf of Chiriquí, and Gulf of Montijo; and in Colombia: the Northern Chocó Region (Gulf of Tribugá, and from Utría National Park north to Juradó). Due to external factors, it was not possible to visit southern areas of the Pacific coast in Colombia, where most of the mangrove forests of the country are distributed. Potential errors associated with this omission are included in the discussion. Land use obtained from land cover maps of different sources (Table A1) was verified via ground-truthing. Table A2 shows the amount of ground-truthed cells from the 1 km² sample grid. Cohen's kappa (κ) was used to determine if there was agreement between land use information (Table A1) and ground-truthing. Results show that there was significant agreement in all countries, with Colombia and Costa Rica showing complete agreement (Colombia $\kappa = 1$, (95% CI, 1 to 1), $p < 0.001$; Costa Rica $\kappa = 1$, (95% CI, 1 to 1), $p < 0.001$), whereas Panamá showed a more moderate but significant agreement (Panamá $\kappa = 0.879$ (95% CI, 0.864 to 0.893), $p < 0.001$; total for 3 countries $\kappa = 0.948$ (95% CI, 0.942 to 0.955), $p < 0.001$).

In Panamá, in some instances, aquaculture was incorrectly classified in the land use maps. In these cases, we manually created maps of aquaculture by delimiting ponds using Google Earth imagery at the best possible resolution calibrated with ground-truthing. Finally, we made sure that all aquaculture was correctly classified at a country level in our maps by verifying them with aquaculture experts from the Aquatic Resources Authority of Panamá (Autoridad de los Recursos Acuáticos de Panamá - ARAP).

Table A2

Total number of 1 km² grid cells of mangroves in the 3 study countries, and the number of cells ground-truthed per country.

Country	Total number of cells	Number of ground-truthed cells
Costa Rica	1225	166
Panamá	4066	244
Colombia	4521	11
Total	9812	401

Table A3

Protected areas of the Pacific coast of Colombia, Costa Rica, and Panamá that include mangroves inside their boundaries. Information on management category, year of creation and area was obtained via www.protectedplanet.net, and when not available, obtained through consultation with the relevant government.

Country	Name of protected area	Management category	Year of creation	Area (km ²)
Colombia	Sanquianga	National Natural Park	1977	866.85
	Utría	National Natural Park	1987	653.67
	Uramba Bahía Málaga	National Natural Park	2010	473.18
	Río Anchicaya	National Protected Forest Reserve	1946	1451.52
	Territorio Colectivo Baudó	Regional District of Integrated Management	2008	68.11
Costa Rica	Parque Natural Regional La Sierpe	Regional Natural Park	2008	252.97
	Caletas-Arío	Wildlife Refuge	2006	204.34
	Cipanci	Wildlife Refuge	2001	34.83
	Corcovado	National Park	1975	444.9
	Estero Puntarenas y manglares	Wetland	2001	51.93

(continued on next page)

Table A3 (continued)

Country	Name of protected area	Management category	Year of creation	Area (km ²)
Panamá	Golfito	Wildlife Refuge	1985	28.19
	Golfo Dulce	Forest Reserve	1978	599.9
	Iguanita	Wildlife Refuge	1994	1.13
	La Ensenada	Wildlife Refuge	1998	4.86
	Las Baulas de Guanacaste	National Park	1991	273.25
	Manglar Terraba-Sierpe	Wetland	1994	261.83
	Manuel Antonio	National Park	1972	1264.64
	Marino Ballena	National Park	1992	53.6
	Ostional	Wildlife Refuge	1983	86.24
	Palo Verde	National Park	1978	172.06
	Palustrino Corral de Piedra	Wetland	1994	24.27
	Pejeperro	Wildlife Refuge	2000	5.95
	Piedras Blancas	National Park	1991	158.43
	Portalón	Wildlife Refuge		2.24
	Playa Hermosa	Wildlife Refuge	1998	27.89
	Río Oro	Wildlife Refuge	1999	17.17
	Santa Rosa	National Park	1966	860.35
	Santuario Ecológico	Wildlife Refuge	2003	3.31
	Tivives	Protected Zone	1986	24.74
	Bahía de Chame	Area of multiple uses	2007	89.00
	Bahía de Panamá	Wetland	2003	489.19
	Patiño	Wetland	1993	131.98
	Golfo de Montijo	Wetland	1990	864.76
	Coiba	National Park	2005	2548.24
	Golfo de Chiriquí	National Park	1994	212.21
	Sarigua	National Park	1984	46.70
	Isla Cañas	Wildlife Refuge	1980	242.85
	La Barqueta	Wildlife Refuge	1994	67.04
	Cenegon del Mangle	Wildlife Refuge	1980	8.43
	Playa Boca Vieja	Wildlife Refuge	1994	35.79
	Pablo Barrio	Wildlife Refuge	2009	150.32
Chepigana	Forest Reserve	1960	363.79	
Canglon	Forest Reserve	1984	286.23	
Filo del Tallo	Hydrological Reserve	1997	122.26	
Isla del Rey	Hydrological Reserve	2006	98.22	
Archipiélago de Las Perlas	Special Zone of Marine and Coastal Management	2007	1601.51	

Table A4

Detailed results for a cumulative model of potential drivers of land use and land cover change adjacent to mangrove forests on the Pacific coasts of Costa Rica, Panamá and Colombia. Figures are calculated as a proportion of the total 1 km² grid cells containing mangroves by country.

Variable	Colombia	Costa Rica	Panamá
	% of cells	% of cells	% of cells
	Cumulative model result		
No potential drivers present	73.4	60.6	42.6
One potential driver present	25.7	32.2	40.9
Two potential drivers present	0.9	6.8	15.7
Three potential drivers present	0	0.4	0.8
	Potential driver of LULCC		
Aquaculture	0	9.8	6.1
Agriculture	25.9	27.8	29.4
Coastal development	1.6	9.4	19.4

Table A5

Detailed results for the cumulative impact model of activities adjacent to mangrove forests inside (in) and outside (out) protected areas of three countries of the Eastern Tropical Pacific. Figures are calculated as a proportion of the total 1km² grid cells containing mangroves by country.

Variable	Colombia		Costa Rica		Panamá	
	% of cells		% of cells		% of cells	
	In	Out	In	Out	In	Out
Cumulative model result						
No potential drivers	92	66.7	68.4	53.2	43.4	42.2
One potential driver present	7.7	32.1	26.8	37.3	41.9	40.3
Two potential drivers present	0.2	0.9	4.7	8.8	13.7	16.7
Three potential drivers present	0	0.2	0.2	0.6	0.9	0.7
Potential driver of LULCC						
Aquaculture	0	0.4	4.7	14.7	6	6.1
Agriculture	8	32.2	23.6	31.7	50.4	18.6
Coastal development	0.2	2	8.4	10.3	15.8	21.3




Fig. A1. Frequency distributions from a cumulative model of potential drivers of land use and land cover change adjacent to mangrove forests on the Pacific coasts of Costa Rica, Panamá and Colombia. Bars show the number of 1km² grid cells corresponding to each category.

References

Alongi, D.M., 2002. Present state and future of the world's mangrove forests. *Environ. Conserv.* 29, 331–349 (Cambridge Univ Press).

Alongi, D.M., 2008. Mangrove forests: resilience, protection from tsunamis, and responses to global climate change. *Estuar. Coast. Shelf Sci.* 76, 1–13.

ANAM, 2004. Programa de Acción Nacional de Lucha contra la Desertificación y Sequía en Panamá. Autoridad Nacional del Ambiente, Panamá.

ANAM, ARAP, 2013. Manglares de Panamá: importancia, mejores prácticas y regulaciones vigentes. Editora Novo Art, S.A., Panamá.

Andam, K.S., Ferraro, P.J., Pfaff, A., Sanchez-Azofeifa, G.A., Robalino, J.A., 2008. Measuring the effectiveness of protected area networks in reducing deforestation. *Proc. Natl. Acad. Sci.* 105, 16089–16094.

Bach, O., 2007. Agricultura e implicaciones ambientales con énfasis en algunas cuencas hidrográficas principales. In: Decimotercer Informe. Estado de la Nación en Desarrollo Humano Sostenible. San José, Costa Rica, pp. 22.

Benfield, S.L., Guzmán, H.M., Mair, J.M., 2005. Temporal mangrove dynamics in relation to coastal development in Pacific Panama. *J. Environ. Manag.* 76, 263–276.

Blanco, J.F., Quiceno, P.A., Jiménez, L.F., Turbay, S., 2011. Estrategias de manejo sostenibles de la actividad pesquera en el PNN Utría. In: Informe final de Investigación, 17 ed. Unidad Administrativa Especial del Sistema de Parques Nacionales Naturales (UAESPENN), Fondo de Patrimonio Natural y Corporación Académica Ambiental de la Universidad de Antioquia, pp. 1–402.

Blanco, J.F., Estrada, E.A., Ortiz, L.F., Urrego, L.E., 2012. Ecosystem-wide impacts of deforestation in mangroves: the Urabá Gulf (Colombian Caribbean) case study. *ISRN Ecol.* 2012, 1–14.

Blanco-Libreros, J., Estrada-Urrea, E., 2015. Mangroves on the edge: Anthrome-dependent fragmentation influences ecological condition (Turbo, Colombia, southern Caribbean). *Diversity* 7, 206–228. <https://doi.org/10.3390/d7030206>.

Bolanos, S.M., 2012. Shrimp Aquaculture and Agudulce: A Broken Partnership. Open

Access Dissertations. pp. 1–169.

Briggs, J.C., 1974. *Marine Zoogeography*. McGraw-Hill, New York.

Bruner, A.G., 2001. Effectiveness of parks in protecting tropical biodiversity. *Science* 291, 125–128.

Cabrera, E., Vargas, D.M., Galindo, G., Garcia, M.C., Ordonez, M.F., Vergara, L.K., Pacheco, A.M., Rubiano, J.C., Giraldo, P., 2011. Memoria técnica de la cuantificación de la deforestación histórica nacional- escalas gruesa y fina. Instituto de Hidrología, Meteorología, y Estudios Ambientales -IDEAM, Bogotá D.C. Colombia, pp. 106.

Chape, S., Harrison, J., Spalding, M., Lysenko, I., 2005. Measuring the extent and effectiveness of protected areas as an indicator for meeting global biodiversity targets. *Philos. Trans. R. Soc. B* 360, 443–455.

Chen, J., Chen, J., Liao, A., Cao, X., Chen, L., Chen, X., et al., 2015. Global land cover mapping at 30m resolution: A POK-based operational approach. *ISPRS J. Photogramm. Remote Sens.* 103, 7–27. <https://doi.org/10.1016/j.isprsjprs.2014.09.002>.

Dahdouh-Guebas, F., Van Pottelbergh, I., Kairo, J.G., Cannicci, S., Koedam, N., 2004. Human-impacted mangroves in Gazi (Kenya): predicting future vegetation based on retrospective remote sensing, social surveys, and tree distribution. *Mar. Ecol. Prog. Ser.* 272, 77–92.

Duke, N.C., et al., 2007. A world without mangroves? *Science* 317, 41b–42b.

Etter, A., McAlpine, C., Wilson, K., Phinn, S., Possingham, H., 2006. Regional patterns of agricultural land use and deforestation in Colombia. *Agric. Ecosyst. Environ.* 114, 369–386.

Fiedler, P.C., Talley, L.D., 2006. Hydrography of the eastern tropical Pacific: a review. *Prog. Oceanogr.* 69, 143–180.

Foley, J.A., et al., 2005. Global consequences of land use. *Science* 309, 570–574.

García, C., 2010. Diagnóstico de las áreas marinas y costeras protegidas, y de las áreas de manejo en el pacífico colombiano. Fundación MarViva, Colombia.

Gaveau, D.L.A., Epting, J., Lyne, O., Linkie, M., Kumara, I., Kanninen, M., Leader Williams, N., 2009. Evaluating whether protected areas reduce tropical deforestation in Sumatra. *J. Biogeogr.* 36, 2165–2175.

- Gibbs, H.K., Ruesch, A.S., Achard, F., Clayton, M.K., Holmgren, P., Ramankutty, N., Foley, J.A., 2010. Tropical forests were the primary sources of new agricultural land in the 1980s and 1990s. *Proc. Natl. Acad. Sci. U. S. A.* 107, 16732–16737.
- Gibson, L., et al., 2011. Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* 478, 378–381.
- Giri, C., Zhu, Z., Tieszen, L.L., Singh, A., Gillette, S., Kelmelis, J.A., 2008. Mangrove forest distributions and dynamics (1975–2005) of the tsunami-affected region of Asia. *J. Biogeogr.* 35, 519–528.
- Giri, C., Ochieng, E., Tieszen, L.L., Zhu, Z., Singh, A., Loveland, T., Masek, J., Duke, N., 2011. Status and distribution of mangrove forests of the world using earth observation satellite data. *Glob. Ecol. Biogeogr.* 20, 154–159.
- Hamilton, S.E., 2013. Assessing the role of commercial aquaculture in displacing mangrove forest. *Bull. Mar. Sci.* 89 (2), 585–601.
- Hamilton, S.E., Casey, D., 2016. Creation of a high spatio-temporal resolution global database of continuous mangrove forest cover for the 21st century (CGMFC-21). *Glob. Ecol. Biogeogr.* 25, 729–738.
- Hansen, M.C., et al., 2013. High-resolution global maps of 21st-century forest cover change. *Science* 342, 850–853.
- Heumann, B.W., 2011. Satellite remote sensing of mangrove forests: recent advances and future opportunities. *Prog. Phys. Geogr.* 35, 87–108.
- Hogarth, P., 2007. *The Biology of Mangroves and Seagrasses*. OUP, Oxford.
- IDEAM, IGAC, IAvH, INVEMAR, Sinchi, IIAP, 2007. *Ecosistemas continentales, costeros y marinos de Colombia*. In: 36 Hojas cartográficas. Bogotá, DC.
- Kaufmann, K., 2012. Nuestros humedales, nuestro futuro. In: Plan de conservación para los humedales de la Bahía de Panamá. Sociedad Audubon de Panamá, Panamá, pp. 73.
- Kothari, A., Cooney, R., Hunter, D., MacKinnon, K., Müller, E., Nelson, F., Oli, K., Pandey, S., Rasheed, T., Vavrova, L., 2015. Managing resource use and development. In: Worboys, G.L., Lockwood, M., Kothari, A., Feary, S., Pulsford, I. (Eds.), *Protected Area Governance and Management*. ANU Press, pp. 789–821.
- Lacerda, L.D., Conde, J.E., Alarcon, C., Alvarez-Leon, R., Bacon, P.R., D'Croz, L., Kjerfve, B., Polaina, J., Vannucci, M., 1993. Mangrove ecosystems of Latin America and the Caribbean: a summary. In: Lacerda, L.D. (Ed.), *Conservation and Sustainable Utilization of Mangrove Forests in Latin America and Africa*. Mangrove Ecosystems Technical Reports International Society for Mangrove Ecosystems, pp. 44.
- Leal, C., 2000. Manglares y economía extractiva. In: Arocha, J., Machado, M.L., Villa, W. (Eds.), *Geografía humana de Colombia: los afrocolombianos*. Tomo VI Instituto Colombiano de Cultura Hispanica, Bogotar.
- López Angarita, J., Moreno-Sánchez, R., Maldonado, J.H., Sánchez, J.A., 2014. Evaluating linked social-ecological systems in marine protected areas. *Conserv. Lett.* 7, 241–252.
- López Angarita, J., Roberts, C.M., Tilley, A., Hawkins, J.P., Cooke, R.G., 2016. Mangroves and people: lessons from a history of use and abuse in four Latin American countries. *For. Ecol. Manag.* 368, 151–162.
- Lucas, R.M., Mitchell, A.L., Rosenqvist, A., Proisy, C., Melius, A., Ticehurst, C., 2007. The potential of L-band SAR for quantifying mangrove characteristics and change: case studies from the tropics. *Aquat. Conserv. Mar. Freshwat. Ecosyst.* 17, 245–264.
- Manson, F.J., Loneragan, N.R., McLeod, I.M., Kenyon, R.A., 2001. Assessing techniques for estimating the extent of mangroves: topographic maps, aerial photographs and Landsat TM images. *Mar. Freshw. Res.* 52, 787–792.
- Mariscal, E., 2012. Causas directas e indirectas de la deforestación y degradación de bosques - cambio de uso de suelo. PNUMA, ANAM, pp. 1–77.
- Miteva, D.A., Murray, B.C., Pattanayak, S.K., 2015. Do protected areas reduce blue carbon emissions? A quasi-experimental evaluation of mangroves in Indonesia. *Ecol. Econ.* 119, 127–135.
- Naughton-Treves, L., Holland, M.B., Brandon, K., 2005. The role of protected areas in conserving biodiversity and sustaining local livelihoods. *Annu. Rev. Environ. Resour.* 30, 219–252.
- Nepstad, D., et al., 2014. Slowing Amazon deforestation through public policy and interventions in beef and soy supply chains. *Science* 344, 1118–1123.
- Nunan, L.M., Arce, S.M., Staha, R.J., Lightner, D.V., 2001. Prevalence of infectious hypodermal and hematopoietic necrosis virus (IHNV) and white spot syndrome virus (WSSV) in *Litopenaeus vannamei* in the Pacific Ocean off the coast of Panama. *J. World Aquacult. Soc.* 32, 330–334.
- Páez-Osuna, F., 2001. The environmental impact of shrimp aquaculture: causes, effects, and mitigating alternatives. *Environ. Manag.* 28, 131–140.
- Parra, A.S., Angel, J.D.R., 2014. El colapso ambiental en el río Patía, Colombia: variaciones morfológicas y alteraciones en los ecosistemas de manglar. *Lat. Am. J. Aquat. Res.* 42 (1), 40–60. <https://doi.org/10.3856/vol42-issue1-fulltext-4>.
- Pinto, I., Yee, J., 2011. Diagnóstico de las áreas marinas protegidas y de las áreas marinas para la pesca responsable en el Pacífico panameño. Fundación MarViva, Panamá.
- Potapov, P.V., Turubanova, S.A., Hansen, M.C., Adusei, B., Broich, M., Altstatt, A., Mane, L., Justice, C.O., 2012. Quantifying forest cover loss in Democratic Republic of the Congo, 2000–2010, with Landsat ETM+ data. *Remote Sens. Environ.* 122, 106–116.
- RAMSAR, 2015. *The Ramsar Convention of Wetlands, 2015 ed.* Available from: http://ramsar.rgis.ch/cda/en/ramsar-documents-wurl-policies-national-wetland-21185/main/ramsar/1-31-116-162%5E21185_4000_0_, Accessed date: 22 June 2017.
- Restrepo, J.D., Cantera, J.R., 2013. Discharge diversion in the Patía River delta, the Colombian Pacific: geomorphic and ecological consequences for mangrove ecosystems. *J. S. Am. Earth Sci.* 46, 183–198. <https://doi.org/10.1016/j.jsames.2011.04.006>.
- Richards, D.R., Friess, D.A., 2015. Rates and drivers of mangrove deforestation in Southeast Asia, 2000–2012. *Proc. Natl. Acad. Sci. U. S. A.* 113, 344–349.
- Salas, E., Ross Salazar, E., Arias, A., 2012. Diagnóstico de las áreas marinas protegidas y áreas marinas para la pesca responsable en el pacífico costarricense. Fundación MarViva, San José, Costa Rica.
- Sánchez-Paez, H., Alvarez-León, R., Guevara-Mancera, O.A., Zamora-Guzmán, A., Rodríguez-Cruz, H., Bravo-Pazmiño, H., 1997. Diagnóstico y zonificación preliminar de los manglares del Pacífico de Colombia. In: Proyecto de conservación y manejo para el uso múltiple y el desarrollo de los manglares de Colombia. Ministerio Del Medio Ambiente, Organización Internacional de Maderas Tropicales, Dirección de Proyectos de Repoblación y Ordenación Forestal.
- Seto, K.C., Fragkias, M., Güneralp, B., Reilly, M.K., 2011. A meta-analysis of global urban land expansion. *PLoS One* 6, e23777.
- Simard, M., Rivera-Monroy, V.H., Mancera-Pineda, J.E., Castañeda-Moya, E., Twilley, R.R., 2008. A systematic method for 3D mapping of mangrove forests based on shuttle radar topography mission elevation data, ICESat/GLAS waveforms and field data: application to Ciénaga Grande de Santa Marta, Colombia. *Remote Sens. Environ.* 112, 2131–2144.
- Spracklen, B.D., Kalamandeen, M., Galbraith, D., Gloor, E., Spracklen, D.V., 2015. A global analysis of deforestation in moist tropical forest protected areas. *PLoS One* 10, e0143886.
- Thomas, N., Lucas, R., Bunting, P., Hardy, A., Rosenqvist, A., Simard, M., 2017. Distribution and drivers of global mangrove forest change, 1996–2010. *PLoS One* 12 (6), e0179302. <https://doi.org/10.1371/journal.pone.0179302>.
- Thompson, I.D., Guariguata, M.R., Okabe, K., 2013. An operational framework for defining and monitoring forest degradation. *Ecol. Soc.* 18, 20.
- Tilman, D., Balzer, C., Hill, J., Befort, B.L., 2011. Global food demand and the sustainable intensification of agriculture. *Proc. Natl. Acad. Sci. U. S. A.* 108, 20260–20264.
- Trejos, N., Morán, M., Smith, O., Morales, M., 2008. Diagnósticos Biofísico, Institucional-Legal, Socioeconómico y Línea Base del Bosque de Manglar del Golfo de Chiriquí, Provincia de Chiriquí. Centro del Agua del Trópico Húmedo para América Latina y el Caribe, pp. 1–134.
- UNEP, 2014. In: van Bochove, J., Sullivan, E., Nakamura, T. (Eds.), *The Importance of Mangroves to People: A Call to Action*. United Nations Environment Programme World Conservation Monitoring Centre, Cambridge.
- Valiela, I., Bowen, J.L., York, J.K., 2001. Mangrove forests: one of the world's threatened major tropical environments. *Bioscience* 51, 807.
- Vieira, C.A., Diaz Granados, M.A., Díaz, J.M., 2016. Ordenamiento y manejo pesquero en la costa norte del Pacífico Chococano. In: Díaz, J.M., Guillot, L., Velandia, C.M. (Eds.), *La pesca artesanal en el norte del Pacífico colombiano: Un horizonte ambivalente*. Fundación MarViva, Bogotá.
- White, E.V., Roy, D.P., 2015. A contemporary decennial examination of changing agricultural field sizes using Landsat time series data. *Geo* 2, 33–54.
- Zapata, F.A., Robertson, R., 2006. How many species of shore fishes are there in the tropical eastern Pacific? *J. Biogeogr.* 34, 38–51.